

***Cordana sinensis* sp. nov. from southern China**

LING QIU¹, KAI ZHANG², ZHAO-HUAN XU¹,
RAFAEL F. CASTAÑEDA-RUÍZ³, JIAN MA^{1,4*}

¹ College of Agronomy, Jiangxi Agricultural University, Nanchang, Jiangxi 330045, China

² College of Forestry Engineering, Shandong Agriculture and Engineering University,
Jinan, Shandong 250100, China

³ Instituto de Investigaciones Fundamentales en Agricultura Tropical ‘Alejandro de Humboldt’
(INIFAT), Académico Titular de la “Academia de Ciencias de Cuba”,
Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

⁴ Jiangxi Key Laboratory for Conservation and Utilization of Fungal Resources,
Jiangxi Agricultural University, Nanchang, Jiangxi 330045, China

* CORRESPONDENCE TO: majian821210@163.com; jxajumj@126.com

ABSTRACT—A new species, *Cordana sinensis*, is described and illustrated from a specimen collected on dead branches of an unidentified broadleaf tree in Jiangxi Province, China. It is characterized by its integrated, polyblastic conidiogenous cells that produce ovoid to obclavate, medially 1–2-septate pale brown, smooth conidia with a slightly prominent hilum. A dichotomous key and a synoptic table to *Cordana* species, are provided.

KEY WORDS—asexual *Ascomycota*, *Cordanaceae*, hyphomycetes, saprobes, taxonomy

Introduction

Cordana Preuss was established with *C. pauciseptata* Preuss as the type species (Preuss 1851) and characterized mainly by macronematous, mononematous, unbranched, cylindrical, nodose or subnodose conidiophores and polyblastic, integrated, terminal or intercalary, sympodial conidiogenous cells, usually denticulate with tiny or conspicuous cylindrical denticles. The conidia are solitary, acropleurogenous, variably shaped (ovoid, obovoid, ellipsoidal, pyriform, or obpyriform), 0–1-euseptate and seceding schizolytically (Preuss 1851, Hughes 1955, Ellis 1971, Markovskaja 2003,

Seifert & al. 2011). Hernández-Restrepo & al. (2014), who expanded the generic concept to include integrate and discrete polyblastic conidiogenous cells that are intercalary nodose and swollen or umbellate at the conidiophore apices, provided a key for 19 accepted species. Thereafter, *C. johnstonii* M.B. Ellis, *C. musae* (Zimm.) Höhn., and *C. versicolor* D.J. Soares & R.W. Barreto were transferred to *Neocordana* Hern.-Rest. & Crous based on a phylogenetic analysis (Hernández-Restrepo & al. 2015). After the addition of four *Cordana* species by Ai & al. (2019) and Luo & al. (2019), *Cordana* currently contains 20 recognized species, of which only ten have been reported from China: *C. abramovii* Seman & Davydkina, *C. aquatica* Z.L. Luo & al., *C. ellipsoidea* de Hoog, *C. lignicola* Z.L. Luo & al., *C. lithuanica* Markovsk., *C. lushanensis* C.C. Ai & al., *C. meilingensis* C.C. Ai & al., *C. pauciseptata*, *C. uniseptata* L. Cai & al., and *C. terrestris* (Timonin) Hern.-Restr. & al. (Lu & al. 2000, Cai & al. 2004, Zhang 2018, Ai & al. 2019, Luo & al. 2019).

In the past, *Cordana* was associated with *Trichosphaeriaceae* (Müller & Samuels 1982) and *Chaetosphaeriaceae* (Réblová & al. 1999) based on ascomatal morphology.

Saprobic hyphomycetes are highly diverse on dead branches, and much taxonomic information has been published in China (e.g., Wu & Zhuang 2005; Ma & al. 2014, 2015, 2016; Ma & Zhang 2015; Xia & al. 2015, 2016; Ma 2016a,b; Xu & al. 2017, 2019; Zhang 2018; Qiu & al. 2020). In our study of these fungi from Jinggangshan Mountain, Jiangxi Province, an interesting hyphomycete with the morphological features typical of *Cordana* was collected on dead branches. It shows significant differences from previously described *Cordana* species and therefore is proposed as new.

Materials & methods

Dead branches were collected from humid environments and watersides in the forest ecosystems of southern China and placed in Ziploc™ plastic bags for transport to the laboratory, where they were processed, and examined as described by Ma & al. (2011). Microphotographs were prepared using a Nikon Eclipse E200 and SmartV550Dc digital camera, with a 100 × (oil immersion) objective at the same background and scale. Conidia were measured at their widest point. The minimum and maximum values for microscopic measurements are provided, with outliers in parentheses. Adobe Photoshop 7.0 was used to assemble photographs into plates. The specimens are deposited in the Herbarium of Jiangxi Agricultural University, Plant Pathology, Nanchang, Jiangxi, China (HJAUP).




FIG. 1. *Cordana sinensis* (holotype, HJAUP M0172).
A–C. Conidiophores and conidia; D. Conidia.

Taxonomy

Cordana sinensis L. Qiu, K. Zhang, R.F. Castañeda & Jian Ma, **sp. nov.**

FIG. 1

MB 836854

Differs from *Cordana crassa* and *C. lithuanica* by its ovoid to obclavate, medially 1-septate or with 2-septate, narrower conidia.

TYPE: China, Jiangxi Province: Jinggangshan Mountain, Jingzhushan scenic spot, 26°32'28"N 114°06'57"E, on dead branches of an unidentified broadleaf tree, 6 Nov. 2014, J. Ma (**holotype**, HJAUP M0172).

ETYMOLOGY: refers to China, where the fungus was collected.

COLONIES on the natural substratum effuse, brown to dark brown. Mycelium partly superficial, partly immersed, composed of branched, septate, smooth, subhyaline to pale brown hyphae. CONIDIOPHORES macronematous, mononematous, simple or branched, erect, straight to flexuous, smooth, septate, brown to pale brown, cylindrical, sub-nodose, ≤ 400 μm long, 4–6.5 μm wide, with terminal and intercalary nodes, 5–7.5 μm wide. CONIDIOGENOUS CELLS integrated, polyblastic, terminal and intercalary, cylindrical-subnodose, with tiny prominent denticles at the nodes, percurrent elongated, $9.5\text{--}32 \times 4\text{--}6.5$ μm . Conidial secession schizolytic. CONIDIA solitary, acropleurogenous, ovoid to obclavate, 1–2-septate, pale brown, smooth, thick-walled, $13\text{--}20(\text{--}26.5) \times 5\text{--}7.5$ μm , dry, apical cell rounded, basal cell with a tiny prominent hilum.

COMMENTS –Among the known species of *Cordana*, *C. sinensis* most closely resembles *C. crassa* Tóth and *C. lithuanica* Markovsk. in conidial shape. However, *C. crassa* and *C. lithuanica* produce asymmetrically 1-septate conidia while those of *C. sinensis* are medially 1–2-septate. In addition, *C. crassa* produces larger (20–24 μm long), ovoid conidia (Tóth 1975), and *C. lithuanica* produces shorter and wider ($13\text{--}17.5 \times 8\text{--}10$ μm) obpyriform conidia (Markovskaja 2003).

With the addition of *C. sinensis*, *Cordana* is represented by 21 species (TABLE 1). All these species have integrated, polyblastic, terminal or intercalary, sympodial conidiogenous cells, except *C. bisbyi* (Timonin) Hern.-Restr. & al. and *C. terrestris* (Timonin) Hern.-Restr. & al., which bear discrete, polyblastic conidiogenous cells umbellately arranged at the apex of the conidiophores (Hernández-Restrepo & al. 2014). *Cordana* conidia are mostly smooth-walled and concolorous, but *C. indica* Subramon. & V.G. Rao and *C. verruculosa* Hern.-Restr. & al. produce echinulate or verruculose conidia. *Cordana inaequalis* S. Hughes, *C. mercadoana* Hern.-Restr. & al.,

TABLE 1 Comparisons of conidia and substrates of *Cordana* species. The new species is in bold.

SPECIES	CONIDIA			HABITAT	TYPE LOCALITY
	SIZE (µm)	SHAPE	SEPTATION	COLOR	
<i>C. abramovi</i> [†]	18–31 × 12.5–15.5	Ovoid	1	Uniform	Fruitbodies of <i>aphyllorphoralean</i> <i>basidiocarp</i> and wood Georgia
<i>C. andinopatagonica</i> [‡]	17–19 × 10–13	Obovoid to ellipsoid	1	Uniform	Leaf litter of <i>Nothofagus dombevi</i> Argentina
<i>C. aquatica</i> [‡]	8–10 × 3–5	Mostly oblong, sometimes ellipsoid	1	Uniform	Submerged decaying wood China
<i>C. bisbyi</i> ^{§,5*}	5–7.2 × 2.5–3.6	Cylindrical–oval	0	Uniform	Soil Honduras
<i>C. crassa</i> [§]	20–24 × 13–15	Ovoid	1	Uniform	Rotting wood of <i>Fagus sylvatica</i> Hungary
<i>C. ellipsoidea</i> [‡]	8.7–11.5 × 4.3–4.7	Ellipsoid to oblong	1	Uniform	Soil Nepal
<i>C. inaequalis</i> [§]	10–12.5 × 4–5	Ellipsoidal to ovoid	1	Variable	Rotten wood Canada
<i>C. indica</i> ^{§,6*}	11–20 × 4.5–7	Oval to oblong	1	Uniform	Dead leaves of <i>Casuarina</i> India
<i>C. lignicola</i> [‡]	9–11 × 3.5–4.5	Mostly oblong, sometimes ellipsoid,	0–2	Uniform	Submerged decaying wood China
<i>C. lithuanica</i> ¹⁰	13–17.5 × 8–10	Obpyriform	1	Uniform	Decaying wood of <i>Fraxinus excelsior</i> Lithuania
<i>C. lushmanensis</i> ¹¹	5.5–8 × 2.5–4	Ellipsoidal to obovoid	0	Uniform	Dead branches China

SPECIES	CONIDIA			HABITAT		TYPE LOCALITY
	SIZE (µm)	SHAPE	SEPTATION	COLOR		
<i>C. meltingensis</i> ¹¹	10–13 × 5.5–7	Oblong or cylindrical	1	Uniform	Dead branches	China
<i>C. mercadoana</i> ⁴	6–10 × 3–4	Oblong, obovoid or cylindrical	0–1	Variable	Dead twig	Spain
<i>C. parasitica</i> ¹²	6–10 × 3–5	Oblong	1	Variable	Living leaves of <i>Aster trinervii</i> var. <i>odusti</i>	Japan
<i>C. pauciseptata</i> ¹³	8–12 × 5–7	Broad ellipsoid to ovoid	1	Uniform	Rotten wood of frondose tree	Germany
<i>C. semaniae</i> ¹⁴	21–27 × 9–15	Obovoid	0	Uniform	Bark of rotten <i>Quercus</i>	Russia
<i>C. sinensis</i>	13–20 × 5–7.5	Ovoid to obclavate	1–2	Uniform	Dead branches	China
<i>C. solitaria</i> ¹⁵	6.5–9.5 × 4.5–6.5	Obovoidal	0	Uniform	Rotten wood	India
<i>C. terrestris</i> ^{4,16,*}	6–11 × 3–4	Oblong to ovate	1	Uniform	Soil	Canada
<i>C. uniseptata</i> ¹⁷	13.5–23 × 8.5–11.5	Mostly broad ellipsoid	1	Variable	<i>Phyllostachys bambusoides</i> submerged in forest stream	China
<i>C. verruculosa</i> ^{4,18**}	3–5.5 × 2–3.5	Ellipsoidal to obovoid	0	Uniform	Dead wood	Spain

*All conidiogenous cells are integrated polyblastic, except for *C. bisbyi* and *C. terrestris*, which are discrete polyblastic.

**All conidia are smooth, except for *C. indica*, which is echinulate and *C. verruculosa*, which is verruculose.

Data from: ¹Seman & Davydkina 1983; ²Gamundi & al. 1979; ³Luo & al. 2019; ⁴Hernández-Restrepo & al. 2014; ⁵Timonin 1961; ⁶Toth 1975;

⁷de Hoog 1973; ⁸Hughes 1983; ⁹Subramoniam & Rao 1976; ¹⁰Markovskaja 2003; ¹¹Ai & al. 2019; ¹²Togashi & Onuma 1934; ¹³Preuss 1851;

¹⁴Davydkina & Mel'nik 1989; ¹⁵Rao & de Hoog 1986; ¹⁶Timonin 1940; ¹⁷Cai & al. 2004.

C. parasitica Togashi & Onuma, and *C. uniseptata* L. Cai & al. produce versicolored conidia. Only five species produce nonseptate conidia: *C. bisbyi*, *C. lushanensis* C.C. Ai & al., *C. semaniae* Davydkina & al., *C. solitaria* V. Rao & de Hoog, and *C. verruculosa*, while most form 1-septate conidia, although in *C. lignicola* Z.L. Luo & al. and *C. sinensis*, 2-septate conidia are occasionally present. All 21 species except *C. parasitica* are usually found on rotten wood or dead branches, or in soil.

Key to *Cordana* accepted species

1. Conidiogenous cells discrete, polyblastic, umbellately arranged over the swollen apex of the conidiophores 2
1. Conidiogenous cells integrated, polyblastic, intercalary or terminal on the conidiophores 3
2. Conidia 0-septate, cylindrical-oval, $5-7.2 \times 2.5-3.6 \mu\text{m}$ *C. bisbyi*
2. Conidia 1-septate, oblong to ovate, $6-11 \times 3-4 \mu\text{m}$ *C. terrestris*
3. Conidia nonseptate 4
3. Conidia septate 7
4. Conidia verruculose, $3-5.5 \times 2-3.5 \mu\text{m}$ *C. verruculosa*
4. Conidia smooth, $\geq 5.5 \mu\text{m}$ long 5
5. Conidia obovoid, guttulate, $21-27 \times 9-15 \mu\text{m}$ *C. semaniae*
5. Conidia $\leq 9.5 \times 6.5 \mu\text{m}$ 6
6. Conidia ellipsoidal to obovoid, $5.5-8 \times 2.5-4 \mu\text{m}$ *C. lushanensis*
6. Conidia obovoidal, $6.5-9.5 \times 4.5-6.5 \mu\text{m}$ *C. solitaria*
7. Conidia oval to oblong, echinulate, $11-20 \times 4.5-7 \mu\text{m}$ *C. indica*
7. Conidia smooth, variably sized 8
8. Conidia versicolorous 9
8. Conidia concolorous 12
9. Conidia ellipsoidal or ovoid 10
9. Conidia oblong, obovoid, or cylindrical 11
10. Conidia mostly broadly ellipsoidal, $13.5-23 \times 8.5-11.5 \mu\text{m}$ *C. uniseptata*
10. Conidia narrowly ellipsoidal to ovoid, $10-12.5 \times 4-5 \mu\text{m}$ *C. inaequalis*
11. Conidia oblong, only 1-septate, $6-10 \times 3-5 \mu\text{m}$ *C. parasitica*
11. Conidia oblong, obovoid or cylindrical, 0-1-septate, $6-10 \times 3-4 \mu\text{m}$ *C. mercadoana*
12. Conidia 0-2 or 1-2-septate 13
12. Conidia only 1-septate 14
13. Conidia mostly oblong, sometimes ellipsoid, 0-2-septate, $9-11 \times 3.5-4.5 \mu\text{m}$ *C. lignicola*
13. Conidia ovoid to obclavate, 1-2-septate, $13-20 \times 5-7.5 \mu\text{m}$ *C. sinensis*

14. Conidia of both cells more or less equally wide 15
14. Conidia of both cells unequally wide 18
15. Conidia ovoid, with a pigmented septal pore,
18–31 × 12.5–15.5 µm *C. abramovii*
15. Conidia oblong, cylindrical, ellipsoid to oblong or ellipsoid to ovoid,
lacking pigmented septal pore, ≤13 × 7 µm 16
16. Conidia ellipsoid to oblong, 8.7–11.5 × 4.3–4.7 µm *C. ellipsoidea*
16. Conidia ≥5 µm wide 17
17. Conidia oblong or cylindrical, 10–13 × 5.5–7 µm *C. meilingensis*
17. Conidia broadly ellipsoid to ovoid, 8–12 × 5–7 µm *C. pauciseptata*
18. Conidia ovoid, 20–24 × 13–15 µm *C. crassa*
18. Conidia obpyriform, obovoid, ellipsoid or oblong, ≤19 × 13 µm 19
19. Conidia mostly oblong, sometimes ellipsoid, 8–10 × 3–5 µm *C. aquatica*
19. Conidia ≥13 × 8 µm 20
20. Conidia obovoid to ellipsoid, 17–19 × 10–13 µm *C. andinopatagonica*
20. Conidia obpyriform, 13–17.5 × 8–10 µm *C. lithuanica*

Acknowledgments

The National Natural Science Foundation of China (Nos. 31970018, 31360011) supported this project. The authors express gratitude to Dr. Rosa María Arias Mota (Instituto Tecnológico Superior de Xalapa, Mexico) and Dr. De-Wei Li (The Connecticut Agricultural Experiment Station, USA) for serving as pre-submission reviewers and to Dr. Lorelei L. Norvell for final editorial review and Dr. Shaun Pennycook for nomenclatural review.

Literature cited

- Ai CC, Ma J, Zhang K, Castañeda-Ruiz RF, Zhang XG. 2019. *Cordana meilingensis* and *C. lushanensis* spp. nov. from Jiangxi, China. Mycotaxon 134: 329–334. <https://doi.org/10.5248/134.329>
- Cai L, McKenzie EHC, Hyde KD. 2004. New species of *Cordana* and *Spadicoides* from decaying bamboo culms in China. Sydowia 56: 222–228.
- Davydkina TA, Mel'nik VA. 1989. Two new hyphomycetes from the genera *Cordana* and *Pyriculariopsis*. Mikologiya i Fitopatologiya 23:110–113.
- de Hoog GS. 1973. A new species of *Cordana* (Dematiaceae, Hyphomycetes). Acta Botanica Neerlandica 22: 209–212.
- Ellis MB. 1971. Dematiaceous hyphomycetes. X. Mycological Papers 125. 30 p.
- Gamundi JJ, Arambarri AM, Bucsinszky AM. 1979. Micoflora de la hojarasca de *Nothofagus dombeii*, II. Darwiniana 22: 189–216.
- Hernández-Restrepo M, Gené J, Mena-Portales J, Cano J, Madrid H, Castaneda-Ruiz RF, Guarro J. 2014. New species of *Cordana* and epitypification of the genus. Mycologia 106: 723–734. <https://doi.org/10.3852/13-122>
- Hernández-Restrepo M, Groenewald JZ, Crous PW. 2015. *Neocordana* gen. nov., the causal organism of *Cordana* leaf spot on banana. Phytotaxa 205: 229–238. <https://doi.org/10.11646/phytotaxa.205.4.2>

- Hughes SJ. 1955. Microfungi I. *Cordana*, *Brachysporium*, *Phragmocephala*. Canadian Journal of Botany 33: 259–268. <https://doi.org/10.1139/b55-019>
- Hughes SJ. 1983. *Cordana inaequalis*. Fungi Canadenses 246. 2 p.
- Lu BS, Hyde KD, Ho WH, Tsui KM, Taylor JE, Wong KM, Yanna, Zhou DQ. 2000. Checklist of Hong Kong fungi. Fungal Diversity Research Series 5. 207 p.
- Luo ZL, Hyde KD, Liu JK, Maharachchikumbura SSN, Jeewon R, Bao DF, Bhat DJ & al. 2019. Freshwater *Sordariomycetes*. Fungal Diversity 99: 451–660. <https://doi.org/10.1007/s13225-019-00438-1>
- Ma J. 2016a. *Sporidesmiella lushanensis* and *S. jiulianshanensis* spp. nov. and a new record from China. Mycotaxon 131: 575–581. <https://doi.org/10.5248/131.575>
- Ma J. 2016b. *Corynesporopsis obclavata* and *Stanjehughesia jiangxiensis* spp. nov. from Lushan Mountain, China. Mycotaxon 131: 583–588. <https://doi.org/10.5248/131.583>
- Ma J, Zhang XG. 2015. A preliminary report of dematiaceous hyphomycetes from dead branches in Jiangxi Province. Biological Disaster Science 38: 290–293.
- Ma J, Wang Y, Ma LG, Zhang YD, Castaneda-Ruiz RF, Zhang XG. 2011. Three new species of *Neosporidesmium* from Hainan, China. Mycological Progress 10: 157–162. <https://doi.org/10.1007/s11557-010-0685-2>
- Ma J, Zhang XG, Castañeda-Ruiz RF. 2014. New species of *Acrodactys* and *Repetophragma* from dead branches in China. Mycotaxon 127: 129–134. <http://doi.org/10.5248/127.129>
- Ma J, Xia JW, Castañeda-Ruiz RF, Zhang XG. 2015. Two new species of *Sporidesmiella* from southern China. Nova Hedwigia 101: 131–137. https://doi.org/10.1127/nova_hedwigia/2015/0254
- Ma J, Zhang XG, Castañeda-Ruiz RF. 2016. *Podosporiopsis*, a new genus of synnematus hyphomycetes from China. Mycotaxon 131: 773–780. <http://doi.org/10.5248/131.773>
- Markovskaja S. 2003. A new species of *Cordana* from Lithuania. Mycotaxon 87: 179–185.
- Müller E, Samuels GJ. 1982. Anamorphs of pyrenomycetous ascomycetes II. *Porosphaerella* gen. nov. and its *Cordana* anamorph. Sydowia 35: 150–155.
- Preuss CGT. 1851. Übersicht untersuchter Pilze, besonders aus der Umgegend von Hoyerswerda. Linnaea 24: 99–153.
- Qiu L, Xu ZH, Shi XG, Wu YX, Kuang WG, Ma J. 2020. Six new *Ellisembia* species in Jiangxi Province. Biological Disaster Science 43: 149–155.
- Rao VG, de Hoog GS. 1986. New or critical hyphomycetes from India. Studies in Mycology 28. 84 p.
- Réblová M, Barr ME, Samuels GJ. 1999. *Chaetosphaeriaceae*, a new family for *Chaetosphaeria* and its relatives. Sydowia 51: 49–70.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.
- Seman EO, Davydkina TA. 1983. De genere *Cordana* Preuss in URSS. Novosti Sistematiki Nizshikh Rastenii 20: 114–118.
- Subramoniam V, Rao VG. 1976. Fungi on garbage from Bombay. Acta Botanica Indica 4(1): 58–62.
- Timonin MI. 1940. The interaction of higher plants and soil micro-organisms. I. Microbial population of rhizosphere of seedlings of certain cultivated plants. Canadian Journal of Research 18: 307–317. <http://doi.org/10.1139/cjr40c-031>
- Timonin MI. 1961. New species of the genus *Pseudobotrytis*. Ceiba 9: 27–29.
- Togashi K, Onuma F. 1934. A list of parasitic fungi collected on Mt. Hayachine, Jwate Prefecture. Bulletin Imperial College of Agriculture and Forestry (Morioka, Japan) 17. 74 p.
- Tóth S. 1975. Some new microscopic Fungi, III. Annales Historico-Natureles Musei Nationalis Hungarici 67: 31–35.

- Wu WP, Zhuang WY. 2005. *Sporidesmium*, *Endophragmiella* and related genera from China. Fungal Diversity Research Series 15. 351 p.
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG. 2015. *Codinaea jianfenglingensis* sp. nov. and new records from southern China. Mycotaxon 130: 835–841. <https://doi.org/10.5248/130.835>
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG. 2016. *Sympodiosynnema*, a new genus of dematiaceous hyphomycetes from southern China. Mycotaxon 131: 45–48. <https://doi.org/10.5248/131.45>
- Xu ZH, Hu DM, Luo YQ, Ma J. 2017. Three species of *Linkosia* and *Spadicoides* new to China. Mycotaxon 132: 243–250. <https://doi.org/10.5248/132.243>
- Xu ZH, Zhang K, Zhang XG, Castañeda-Ruiz RF, Ma J. 2019. *Cacumisporium fusiforme* sp. nov. from Jiangxi, China. Mycotaxon 134: 275–279. <https://doi.org/10.5248/134.275>
- Zhang XG. 2018. Flora fungorum sinicorum. Vol. 55. *Corynespora* et genera cognata. Science Press, Beijing.